Alabama Medicaid Pharmacy Morphine Milligram Equivalents (MME) Cumulative Daily Override

FAX: (800) 748-0116 Phone: (800) 748-0130 Fax or Mail to Kepro P.O. Box 3210 Auburn, AL 36831-3210

	PATIENT INFORMATION		
Patient Name	Patient Medicaid #		
Patient DOB	Patient phone # with area code		
	PRESCRIBER INFORMATION		
Prescriber name	NPI #License #		
Phone # with area code	Fax # with area code		
Address (Optional)			
that this treatment is indicate	ged the patient cash for this office visit or for the treatment of this patient's pain mai ted and necessary and meets the guidelines for use as outlined by the Alabama Med I within this request is accurate. I will be supervising the patient's treatment. Suppor rd.	icaid Age	ency. I attest
	Prescribing Provider Signature Date	•	
	DRUG/CLINICAL INFORMATION		
Drug Requested	StrengthDrug Code		
Quantity Requested	Days' Supply for Quantity Requested Date of Last Urine Drug S	creen	
Diagnosis/Specific Descrip	tion of Pain/Medical Justification		
mutual responsibility	for which tapering is not recommended, submit a pain care agreement/conties (with an emphasis on functional improvement). S Below (check yes or no): viewed the patient's PDMP within the past 30 days prior to prescribing the no?	Yes	No No
Has the patient been on aloxone*?	educated on being a candidate to carry naloxone* and/or prescribed		
	nas the patient been counseled on the risk of being/ becoming pregnant of medication, including the risk of neonatal abstinence syndrome (NAS)?		
	unseled the patient on the risk of concurrent use of the requested odiazepines, sedative/hypnotics, or barbiturates?		
Does the patient curre asthma, or hypercarb	ently suffer from respiratory depression, acute or severe bronchial ia?		
^General pain diagnoses will r	not be approved		
should evaluate risk factors fo including considering offering use disorder, higher opioid do:	scribing Opioids for Chronic Pain, before starting and periodically during continuation of oping ropioid-related harms. Clinicians should incorporate into the management plan strategies naloxone when factors that increase risk for opioid overdose, such as history of overdose, sages (≥50 MME/day), or concurrent benzodiazepine use, are present. //65/rr/rr6501e1.htm?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fmmwr%2Fvolumes%2F65%2Frr	to mitigate history of	e risk, substance
	DISPENSING PHARMACY INFORMATION May Be Completed by Pharmacy		
Dispensing pharmacy	NPI #		
	Fax # with area code		
		· <u> </u>	_

Form 413 Effective 12/1/21